



Prevention Research Center for Healthy Neighborhoods
at Case Western Reserve University

2017 Cuyahoga County Youth Risk Behavior Survey: Regional Prevalence

Introduction

The Prevention Research Center for Healthy Neighborhoods (PRCHN) at Case Western Reserve University has administered the Youth Risk Behavior Survey (YRBS) in school districts throughout Cuyahoga County since 2000. The YRBS is a cross-sectional tool developed by the Centers for Disease Control and Prevention (CDC) to track adolescent risk behavior over time. In spring of 2017, the PRCHN conducted the YRBS among 9th through 12th grade students in Cuyahoga County high schools. A more detailed description of the methodology is available at our website, <http://www.prchn.org/YRBSDataCollectionAnalysis.aspx>.

Regional Prevalence


This brief report presents results from the 2017 Cuyahoga County High School Youth Risk Behavior Survey and allows for comparison of health risk behaviors by geographic region. In this brief report, we present:

- [Behaviors that Contribute to Unintentional Injuries](#)
- [Violence](#)
- [Depressive Symptoms and Suicide](#)
- [Tobacco and Electronic Vapor Use](#)
- [Alcohol Use](#)
- [Illegal and Prescription Drug Use](#)
- [Sexual Behaviors](#)
- [Obesity and Weight Control](#)
- [Dietary Behavior](#)
- [Physical Activity](#)
- [Positive Youth Development](#)
- [Preventive Health Care](#)
- [Other Health Behaviors](#)

The overall response rate for inner and outer ring east (57% and 52%, respectively) **did not** reach the standard of 60%, which allows for the data to be considered representative of all students, even those that did not complete the survey. Since the overall response rate was close to the standard of 60%, we have included weighted estimates for these regions. A weight was applied to each record to adjust for student non response and the distribution of students by grade, gender, race/ethnicity, and geographic region within Cuyahoga County. The stability of the data should be considered, given the lower than required overall response rate, and results should be interpreted with caution. In addition, the overall response rate for outer ring west was insufficient for analysis.

Defining the Geographic Regions

The map below provides a visual representation of the six regions of Cuyahoga County. The division of Cuyahoga County into six regions allows for a geographic examination of the prevalence of different risk behaviors in Cuyahoga County. The six regions are:



- Cleveland/CMSD—East: the portion of the Cleveland Metropolitan School District located east of the Cuyahoga River
- Cleveland/CMSD—West: the portion of the Cleveland Metropolitan School District located west of the Cuyahoga River
- Inner Ring—East: a city on the east side of the Cuyahoga River in Cuyahoga County which shares a border with the city of Cleveland
- Inner Ring—West: a city on the west side of the Cuyahoga River in Cuyahoga County which shares a border with the city of Cleveland
- Outer Ring—East: a city on the east side of the Cuyahoga River in Cuyahoga County which does not share a border with the city of Cleveland, and
- Outer Ring—West: a city on the west side of the Cuyahoga River in Cuyahoga County which does not share a border with the city of Cleveland.

Behaviors that Contribute to Unintentional Injuries

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Rarely or never wore a bicycle helmet (Among students who had ridden a bicycle during the 12 months before the survey.)	95.2 (94.1-96.1)	94.2 (93.0-95.1)	88.8 (86.6-90.7)	89.3 (85.3-92.3)	70.2 (67.5-72.7)	n/a
Rarely or never wore a seatbelt (When riding in a car driven by someone else.)	20.5 (18.7-22.4)	18.4 (16.5-20.5)	15.0 (12.5-17.8)	8.5 (7.0-10.2)	8.1 (6.8-9.6)	n/a
Rode with a driver who had been drinking alcohol (During the past 30 days.)	29.8 (27.8-31.8)	26.5 (24.4-28.7)	22.9 (20.6-25.3)	19.8 (17.1-22.8)	17.4 (15.4-19.7)	n/a
Drove a car or other vehicle when student had been drinking alcohol (During the 30 days before the survey.)	9.8 (8.0-12.0)	10.7 (8.9-12.8)	9.3 (7.5-11.4)	9.2 (6.9-12.2)	4.1 (2.6-6.4)	n/a
Texted or e-mailed while driving (Among students who had driven a vehicle during the 30 days before the survey.)	25.1 (22.1-28.4)	22.8 (20.2-25.7)	32.9 (28.1-38.2)	37.1 (33.6-40.7)	42.4 (37.8-47.1)	n/a

Violence

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Carried a weapon (Such as a gun, knife, or club; one or more times during the 30 days before the survey.)	18.4 (16.7-20.2)	19.7 (17.8-21.7)	13.1 (11.0-15.6)	15.4 (13.5-17.5)	11.1 (9.4-13.2)	n/a
Could get and be ready to fire a loaded gun (In less than 24 hours.)	42.4 (40.0-44.8)	42.7 (40.2-45.2)	38.0 (34.5-41.6)	46.5 (43.4-49.7)	37.9 (35.0-40.9)	n/a
Did not go to school because of safety concerns (One or more times during the 30 days before the survey.)	12.0 (10.5-13.7)	12.0 (10.5-13.6)	13.6 (11.6-15.9)	10.9 (9.4-12.6)	12.4 (10.7-14.3)	n/a
Threatened or injured with a weapon, on school property (Such as a gun, knife, or club; one or more times during the 12 months before the survey.)	12.2 (10.7-13.8)	11.6 (10.2-13.3)	9.6 (7.8-11.8)	9.4 (8.1-10.8)	7.1 (5.3-9.4)	n/a
In a physical fight (One or more times during the 12 months before the survey.)	39.1 (36.7-41.6)	34.0 (31.6-36.5)	33.9 (31.2-36.7)	21.1 (18.5-23.9)	20.9 (18.7-23.3)	n/a
Forced to have sexual intercourse	9.1 (8.1-10.2)	10.8 (9.4-12.4)	7.8 (6.3-9.6)	7.2 (6.0-8.6)	3.7 (3.0-4.5)	n/a

Violence

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Forced to do sexual things by someone they were dating or going out with (Such as kissing, touching, or being physically forced to have sexual intercourse; one or more times during the 12 months before the survey, among the students dating.)	11.2 (9.9-12.6)	13.3 (11.3-15.5)	11.3 (9.4-13.7)	13.5 (11.6-15.6)	11.4 (9.4-13.8)	n/a
Physically injured by someone they were dating or going out with (Such as being hit, slammed into something, or injured with an object or weapon; one or more times during the 12 months before the survey, among the students dating.)	13.1 (11.5-14.9)	12.7 (10.7-14.9)	10.7 (9.0-12.7)	9.3 (7.8-11.1)	9.0 (7.4-10.9)	n/a
Bullied on school property (During the 12 months before the survey.)	12.0 (10.7-13.4)	16.9 (15.0-19.0)	12.5 (11.0-14.0)	22.5 (20.6-24.6)	18.1 (15.5-21.0)	n/a
Electronically bullied (Such as through e-mail, chat rooms, instant messaging, websites, or text messaging; during the 12 months before the survey.)	10.0 (8.8-11.3)	14.2 (12.6-15.9)	11.0 (9.2-13.1)	17.2 (15.6-19.0)	13.8 (11.8-16.0)	n/a

Depressive Symptoms and Suicide

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Purposely hurt self without wanting to die (Such as cutting or burning self; during the 12 months before survey.)	20.3 (18.6-22.2)	22.2 (20.4-24.0)	15.7 (14.3-17.2)	21.9 (19.7-24.3)	11.7 (10.1-13.6)	n/a
Felt sad or hopeless (Almost every day for 2 weeks or more in a row; during the 12 months before the survey.)	32.3 (30.2-34.4)	36.6 (34.2-39.0)	30.5 (28.8-32.3)	37.4 (34.8-40.0)	23.5 (21.0-26.2)	n/a
Seriously considered attempting suicide (During the 12 months before the survey.)	16.8 (15.3-18.4)	18.9 (17.1-20.9)	17.6 (15.7-19.7)	20.7 (18.4-23.1)	16.4 (13.1-20.3)	n/a
Attempted suicide (One or more times during the 12 months before the survey.)	17.6 (15.9-19.4)	19.0 (17.2-20.9)	13.9 (11.7-16.4)	10.5 (9.1-12.0)	7.8 (5.4-11.1)	n/a
Knew someone who completed suicide	24.1 (22.2-26.2)	32.6 (30.5-34.9)	24.1 (21.8-26.6)	34.1 (31.9-36.3)	24.9 (22.5-27.4)	n/a

Tobacco and Electronic Vapor Use

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever smoked cigarettes (Smoked a cigarette, even 1 or 2 puffs, on at least 1 day during their life.)	16.1 (14.6-17.7)	25.2 (23.0-27.6)	14.2 (12.4-16.2)	28.1 (24.9-31.5)	15.6 (12.9-18.7)	n/a
Current cigarette use (Smoked a cigarette on at least 1 day during the 30 days before the survey.)	2.7 (2.0-3.5)	6.7 (5.7-7.9)	2.7 (1.9-3.9)	10.6 (8.6-13.0)	4.4 (3.0-6.4)	n/a
Smoked more than 10 cigarettes (Per day, on the days smoked; among the students who smoked 30 days before the survey.)	7.5 (4.5-12.0)	8.0 (4.7-13.2)	12.0 (7.0-19.9)	5.4 (3.3-8.9)	23.2 (7.9-51.4)	n/a
Current cigar use (Smoked a cigar, cigarillo, little cigar, or flavored cigar on at least 1 day during the 30 days before the survey.)	16.8 (15.3-18.5)	16.7 (14.5-19.1)	15.1 (12.8-17.8)	14.5 (12.0-17.4)	10.3 (8.6-12.3)	n/a
Smoked more than 10 cigars, cigarillos, little cigars, or flavored cigars per day (On the days smoked; among the students who smoked 30 days before the survey.)	6.0 (4.2-8.4)	9.3 (6.4-13.4)	9.4 (6.7-12.9)	4.5 (2.6-7.4)	2.4 (2.0-3.0)	n/a
Current use of tobacco in a waterpipe (Smoked from a hookah, narghile or water pipe on at least 1 day during the 30 days before the survey.)	6.5 (5.4-7.8)	10.5 (8.9-12.4)	7.1 (5.6-9.1)	9.3 (7.7-11.3)	7.2 (5.7-9.0)	n/a
Tried first tobacco product before age 13 years (Including cigarettes, cigars, little cigars, flavored cigars, shisha or hookah tobacco, and chewing tobacco.)	17.7 (16.1-19.5)	18.4 (16.4-20.7)	13.6 (11.2-16.5)	10.9 (9.4-12.5)	5.2 (4.4-6.2)	n/a

Tobacco and Electronic Vapor Use

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever tried to quit using all tobacco products (Among students who smoked during the 12 months before the survey.)	45.1 (41.6-48.5)	49.4 (45.8-53.0)	44.3 (38.4-50.4)	45.6 (40.4-50.9)	36.1 (33.2-39.0)	n/a
Ever used an electronic vapor product	25.1 (23.2-27.2)	41.1 (38.5-43.8)	27.8 (25.2-30.5)	40.8 (37.3-44.3)	37.9 (34.1-41.8)	n/a
Current electronic vapor product use (Used at least 1 day during the 30 days before the survey.)	8.0 (6.9-9.1)	16.3 (14.7-18.0)	9.0 (7.4-11.0)	17.7 (15.2-20.5)	19.8 (17.4-22.5)	n/a

Alcohol Use

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever used alcohol (Had at least 1 drink of alcohol on at least 1 day during their life.)	54.2 (51.5-56.9)	59.5 (56.9-62.0)	56.9 (53.2-60.6)	63.2 (60.0-66.2)	56.4 (52.6-60.2)	n/a
Drank alcohol before age 13 years (Other than a few sips.)	19.4 (17.4-21.5)	21.5 (19.1-24.0)	17.6 (14.7-20.9)	15.8 (13.9-17.9)	10.5 (8.0-13.6)	n/a
Current alcohol use (Had at least 1 drink of alcohol on at least 1 day during the 30 days before the survey.)	23.7 (21.5-26.0)	33.3 (30.6-36.2)	26.1 (23.1-29.3)	34.0 (29.8-38.5)	30.7 (27.6-33.9)	n/a
Had 4 or more drinks in a row (if female) or 5 or more drinks (if male) within a couple of hours (During 30 days before the survey.)	12.8 (11.1-14.7)	19.1 (17.1-21.3)	11.6 (9.3-14.4)	18.4 (15.2-22.2)	16.0 (13.7-18.7)	n/a

Illegal and Prescription Drug Use

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever used marijuana (Used marijuana 1 or more times during their life.)	47.2 (44.2-50.3)	44.3 (41.4-47.2)	47.0 (43.5-50.5)	39.4 (34.6-44.4)	36.0 (33.4-38.8)	n/a
Tried marijuana before age 13 years	14.7 (13.0-16.6)	15.5 (13.2-18.1)	10.6 (8.4-13.4)	5.7 (4.5-7.3)	5.0 (4.0-6.2)	n/a
Current marijuana use (Used marijuana 1 or more times during the 30 days before the survey.)	28.1 (25.4-31.0)	25.6 (23.1-28.3)	30.1 (27.2-33.2)	24.7 (21.2-28.6)	22.7 (20.5-25.1)	n/a
Usually smoked marijuana in a blunt (Among students who used marijuana during the 30 days before the survey.)	83.2 (79.3-86.5)	71.0 (66.5-75.0)	73.3 (70.0-76.4)	53.5 (48.8-58.1)	47.2 (42.3-52.1)	n/a
Ever used prescription pain medicine without a doctor's prescription or differently than how a doctor prescribed (Including all or some of the following: codeine, Vicodin, OxyContin, Hydrocodone, or Percocet; one or more times during their life.)	16.8 (15.1-18.7)	19.5 (17.3-21.8)	15.6 (13.2-18.4)	14.3 (11.6-17.4)	7.6 (6.2-9.4)	n/a

Illegal and Prescription Drug Use

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever used synthetic drugs (Including all or some of the following: “designer drugs” such as synthetic marijuana, K2, Spice, fake weed, King Kong, Yucatan Fire, Skunk, Moon Rocks, herbal incense, or bath salts; one or more times during their life.)	8.9 (7.3-10.7)	11.6 (9.8-13.8)	8.6 (6.7-11.0)	6.5 (5.1-8.3)	5.1 (3.1-8.3)	n/a
Ever used an illicit drug (Including all or some of the following: any form of cocaine, heroin, methamphetamines, ketamine, speed, LSD, or ecstasy; one or more times during their life.)	7.2 (5.7-9.1)	9.4 (7.8-11.4)	7.0 (5.6-8.8)	7.8 (6.3-9.7)	4.0 (2.0-7.9)	n/a
Offered, sold, or given illegal drugs on school property (One or more times during the 12 months before the survey.)	17.8 (16.1-19.6)	19.0 (17.1-21.0)	22.7 (20.4-25.2)	16.6 (14.5-18.8)	17.5 (14.8-20.6)	n/a

Sexual Behaviors

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever had sexual intercourse (Had sexual intercourse 1 or more times during their life.)	53.1 (49.9-56.2)	45.5 (41.7-49.5)	48.3 (44.9-51.7)	38.8 (34.8-43.0)	36.8 (34.0-39.8)	n/a
Had sexual intercourse before age 13 years	11.5 (9.8-13.4)	7.8 (6.4-9.5)	7.6 (5.7-10.0)	2.8 (2.1-3.6)	4.1 (2.7-6.2)	n/a
Had sexual intercourse with 4 or more people (Among students who had ever had sexual intercourse.)	16.4 (14.5-18.5)	12.2 (10.4-14.3)	14.0 (11.4-17.0)	6.3 (4.6-8.6)	7.9 (6.8-9.1)	n/a
Currently sexually active (Had sexual intercourse with at least 1 person during the 3 months before the survey.)	35.4 (32.5-38.5)	30.9 (27.4-34.7)	33.5 (30.4-36.8)	27.1 (23.9-30.6)	27.6 (24.9-30.6)	n/a
Drank alcohol or used drugs before having sexual intercourse (Among students who were currently sexually active.)	15.1 (12.5-18.1)	18.0 (14.6-22.1)	15.5 (12.0-19.8)	18.3 (12.9-25.2)	21.6 (18.8-24.7)	n/a
Used a condom during last sexual intercourse (Among students who were currently sexually active.)	51.8 (48.0-55.5)	45.9 (42.2-49.6)	49.3 (44.1-54.6)	60.2 (53.8-66.3)	69.7 (66.2-72.9)	n/a
Been pregnant or gotten someone pregnant (One or more times during their life.)	6.5 (5.4-7.8)	4.5 (3.5-5.7)	4.5 (3.2-6.2)	1.9 (1.4-2.6)	1.5 (1.0-2.2)	n/a

Sexual Behaviors

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ever been taught about AIDS or HIV infection in school	69.3 (66.9-71.7)	65.4 (62.4-68.2)	79.3 (75.7-82.5)	84.5 (82.4-86.4)	85.3 (81.8-88.3)	n/a
Ever been tested for HIV, the virus that causes AIDS (Not including tests done if donated blood.)	23.8 (21.5-26.3)	21.9 (19.9-24.0)	16.3 (13.7-19.2)	9.0 (7.6-10.7)	9.9 (8.7-11.2)	n/a
Adults in family have talked about expectations with students about sex (What to do or not to do about sex.)	69.5 (67.1-71.8)	60.1 (57.7-62.5)	66.1 (62.1-69.8)	55.2 (52.7-57.6)	60.4 (57.9-62.8)	n/a

Obesity and Weight Control

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Obese (Having a BMI of >95 th percentile for age and sex. Calculated from self-reported height and weight, adjusted for sex and age.)	21.8 (20.2-23.4)	21.1 (18.9-23.5)	14.3 (12.8-15.9)	17.3 (15.6-19.1)	13.1 (10.5-16.2)	n/a
Overweight (Having a BMI >85 th percentile and <95 th percentile for age and sex. Calculated from self-reported height and weight, adjusted for sex and age.)	19.4 (17.8-21.2)	17.7 (15.7-19.8)	20.7 (18.0-23.7)	17.2 (15.2-19.5)	16.3 (14.1-18.6)	n/a
Describes self as slightly or very overweight	30.9 (28.9-32.9)	30.5 (28.1-33.0)	28.5 (26.3-30.9)	35.4 (32.7-38.1)	29.1 (25.8-32.5)	n/a
Trying to lose weight	42.9 (40.6-45.3)	44.3 (42.4-46.3)	40.9 (38.2-43.7)	46.6 (44.0-49.2)	42.1 (39.5-44.7)	n/a

Dietary Behavior

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Ate fruits and vegetables 5 or more times per day (During the 7 days before the survey.)	18.2 (16.5-20.0)	20.8 (18.7-23.0)	19.1 (17.0-21.3)	15.2 (13.7-16.8)	24.9 (23.0-26.9)	n/a
Drank water at school on average every day (Including water from a fountain, a sink or faucet, a bottle, a reusable bottle student brought from home, or some other source; during an average school week.)	47.1 (44.7-49.5)	43.4 (40.7-46.2)	51.9 (47.6-56.1)	57.0 (53.2-60.8)	62.5 (59.3-65.5)	n/a
Did not eat breakfast every day (During the 7 days before the survey.)	81.9 (80.1-83.6)	80.9 (79.1-82.7)	78.4 (75.3-81.1)	73.9 (71.7-76.0)	63.4 (60.2-66.4)	n/a
Ate fast food meal or snack 1 or more days (Includes McDonalds, Taco Bell, or KFC; during the 7 days before the survey.)	77.2 (75.3-78.9)	74.2 (72.1-76.2)	74.9 (72.2-77.4)	76.7 (74.3-79.0)	60.6 (57.5-63.7)	n/a
Went hungry because not enough food in the home (Sometimes, most of the time, or always; during the 30 days before the survey.)	20.2 (18.2-22.3)	21.1 (19.4-23.0)	21.2 (18.3-24.4)	19.1 (17.0-21.4)	14.8 (12.7-17.2)	n/a

Physical Activity

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Participated in 60 or more minutes of physical activity on 5 or more days (During the 7 days before the survey.)	27.1 (24.5-29.9)	28.5 (26.0-31.0)	38.7 (35.7-41.7)	40.7 (37.7-43.7)	47.3 (44.3-50.3)	n/a
Did not participate in 60 or more minutes of physical activity on any day (During the 7 days before the survey.)	26.2 (24.3-28.3)	26.5 (24.5-28.6)	21.3 (18.7-24.2)	14.3 (12.8-15.9)	15.7 (14.2-17.4)	n/a
Watched television 3 or more hours per day (On an average school day.)	33.6 (31.6-35.6)	30.2 (28.0-32.4)	30.6 (27.9-33.4)	17.7 (15.7-19.9)	23.4 (20.5-26.6)	n/a
*Played video or computer games or used a computer 3 or more hours per day (On average school day for something that was not school work; includes use of Xbox, PlayStation, iPad or other tablet, smartphone, YouTube, Instagram, Facebook, or other social media.)	48.4 (46.3-50.5)	47.7 (45.5-50.0)	44.8 (41.4-48.3)	48.2 (46.0-50.3)	42.0 (39.2-44.9)	n/a
Played on 1 or more sports teams (During the past 12 months before the survey.)	53.7 (51.0-56.4)	47.0 (44.4-49.7)	57.6 (54.9-60.2)	57.0 (54.1-59.8)	67.9 (63.6-71.8)	n/a
Walk or ride bike to or from school 3 or more days (During an average school week when the weather allows.)	30.0 (27.8-32.3)	30.6 (28.7-32.7)	34.9 (30.7-39.4)	37.9 (34.6-41.4)	10.0 (8.9-11.2)	n/a

Positive Youth Development

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Spent at least 1 day in clubs or organizations outside of school (During the 7 days before the survey.)	46.8 (44.3-49.3)	39.4 (36.6-42.2)	49.5 (47.0-51.9)	42.3 (39.9-44.8)	52.7 (49.5-55.8)	n/a
Spent at least 1 hour helping other people without getting paid to make one's community a better place for people to live (During the 7 days before the survey.)	41.9 (39.5-44.4)	38.9 (36.1-41.7)	37.1 (34.5-39.8)	37.8 (35.4-40.2)	38.8 (36.2-41.6)	n/a
Parents talk with student about school almost every day	48.2 (46.0-50.3)	43.1 (41.1-45.2)	46.3 (43.1-49.5)	47.1 (44.8-49.5)	57.4 (54.6-60.1)	n/a
Students help decide what goes on in school (Agree and strongly agree.)	48.4 (45.9-50.8)	49.7 (47.2-52.2)	49.9 (47.9-52.0)	48.9 (46.0-51.9)	46.5 (44.4-48.6)	n/a
Students feel like they matter to people in their community (Agree and strongly agree.)	43.4 (40.8-46.0)	40.4 (38.3-42.5)	42.3 (38.3-46.4)	39.6 (37.0-42.2)	45.4 (41.3-49.5)	n/a
Obtained 8 or more hours of sleep (On an average school night.)	20.2 (18.1-22.5)	21.1 (19.0-23.3)	18.8 (16.6-21.3)	20.9 (19.1-22.8)	20.6 (18.3-23.1)	n/a
Had a supportive adult (One or more adults, other than a parent, who student would feel comfortable seeking help from if there was an important issue or question affecting their life.)	76.9 (74.9-78.9)	70.6 (68.2-72.9)	76.9 (74.6-79.0)	75.3 (72.9-77.5)	80.9 (78.3-83.2)	n/a
Described their grades in school as A's and B's (During the 12 months before the survey.)	68.3 (65.6-70.9)	65.7 (62.8-68.6)	73.3 (69.9-76.4)	66.1 (61.7-70.1)	84.2 (82.3-85.8)	n/a

Preventive Health Care

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Described health in general as fair or poor	21.0 (19.1-23.1)	23.7 (21.7-25.9)	18.9 (16.6-21.3)	21.2 (18.6-24.1)	17.1 (14.0-20.6)	n/a
Saw a doctor or nurse for check-up or physical exam (During the 12 months before the survey.)	62.3 (59.5-64.9)	58.9 (56.5-61.2)	68.9 (64.5-73.0)	68.8 (66.0-71.5)	77.4 (74.6-80.0)	n/a
Saw a doctor, nurse, or counselor about stress, depression, or problems with emotions (During the 12 months before the survey.)	23.4 (21.6-25.3)	26.2 (24.0-28.6)	21.5 (19.2-23.9)	24.0 (21.7-26.4)	21.6 (18.6-24.9)	n/a

Other Health Behaviors

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
Lives with 2 parents (Biological or Step-parent)	34.6 (32.5-36.8)	48.5 (46.5-50.6)	46.4 (43.3-49.6)	62.4 (59.7-65.1)	67.7 (65.0-70.2)	n/a
Never or rarely feels safe and secure in neighborhood	28.8 (26.4-31.2)	21.7 (19.6-24.1)	13.6 (10.9-16.7)	6.3 (4.8-8.1)	4.9 (2.8-8.6)	n/a
Parent(s) or guardian(s) have been in prison or jail (During the 12 months before the survey.)	15.7 (13.9-17.7)	16.3 (14.5-18.3)	13.4 (11.5-15.6)	6.6 (5.6-7.7)	5.9 (5.0-7.0)	n/a
Had been stopped, questioned, or searched by police (One or more times during the 12 months before the survey.)	24.6 (22.4-26.9)	23.6 (21.0-26.4)	25.6 (22.5-29.0)	23.2 (20.8-25.8)	24.1 (22.2-26.1)	n/a
Described encounters with police as negative or mostly negative (Of those who had encounters with the police during the 12 months before the survey.)	32.2 (29.0-35.5)	31.3 (27.7-35.2)	35.7 (29.7-42.3)	23.5 (20.9-26.3)	24.8 (22.7-27.1)	n/a

Contact Information

Prepared by:

Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
10900 Euclid Avenue
Cleveland, Ohio 44106-7069
www.prchn.org

For more information please contact:

Erika S. Trapl, PhD
Assistant Professor, Department of Population & Quantitative Health Sciences
Associate Director, Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
Erika.trapl@case.edu
216-368-0098

Jean L. Frank, MPH
Manager of School-based Surveillance and Evaluation
Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
Jean.frank@case.edu
216-368-5913

Audrey S. Kinsella, MPH
Data Coordinator
Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
Audrey.kinsella@case.edu
216-368-5746

Briana McIntosh, MPH, CPH
YRBS School-based Surveillance Field Coordinator
Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
Briana.Mcintosh@case.edu
216-368-6571

Marisa Hollinshead, BS
Graduate Assistant
Prevention Research Center for Healthy Neighborhoods
Case Western Reserve University
Marisa.Hollinshead@case.edu

Funding

This work is a product of a Health Promotion and Disease Prevention Research Center supported by Cooperative Agreement Number 1U48DP005030 from the Centers for Disease Control and Prevention and funded either in whole or in part by Cooperative Agreement #U87PS004165-02 from the U.S. Centers for Disease Control and Prevention, Division of Adolescent School Health (CDC-DASH). The findings and conclusions in this report are those of the author(s) and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

CWRU IRB Protocol #2012-274.

