



# 2014 Cuyahoga County Youth Risk Behavior Survey: Tobacco and Electronic Vapor Use

## Introduction

---

The Prevention Research Center for Healthy Neighborhoods (PRCHN) at Case Western Reserve University has administered the Youth Risk Behavior Survey (YRBS) in school districts throughout Cuyahoga County since 2000. The YRBS is a cross-sectional tool developed by the Centers for Disease Control and Prevention (CDC) to track adolescent risk behavior over time. In spring of 2016, the PRCHN conducted the YRBS among 7<sup>th</sup> and 8<sup>th</sup> grade students in Cuyahoga County middle schools. A more detailed description of the methodology is available at our website, <http://prchn.org/yrbs.aspx>.

This brief report presents results from the 2016 Cuyahoga County Middle School Youth Risk Behavior Survey, with a particular focus on Tobacco Use. This includes questions pertaining to tobacco use behavior as well as perceptions of tobacco use. In this brief report, we present:

- [Overall Prevalence](#)
- [Regional Prevalence](#)
- [Demographic Prevalence](#)
- [Trend Report](#)

Data for similar behaviors can be found at our website, <http://prchn.org/yrbs.aspx>.

## Tobacco Use

---

Using tobacco can have serious effects on long-term health. The use of cigarettes is the single leading preventable cause of death in the United States.<sup>i</sup> Almost 90% of adult smokers initiate use before or at age 18.<sup>ii</sup> Hookah smoking can lead to cardiovascular disease and cancer. Per a report from the World Health Organization (WHO), one hookah smoking session may be equivalent to smoking 100 cigarettes.<sup>iii</sup> Tobacco use in adolescence is associated with many other health risk behaviors, including higher-risk sexual behavior and use of alcohol or other drugs.<sup>iv</sup>

## Overall Prevalence

The following table summarizes tobacco use among the 13,261 middle school students who completed the 2016 Cuyahoga County YRBS. Prevalence estimates and 95% confidence intervals were computed for all variables.

Risk Behavior	% (95% Confidence Interval)
<b>Ever smoked cigarettes</b> (Smoked a cigarette, even one or two puffs, on at least 1 day during their life.)	7.8 (7.4-8.2)
<b>Current cigarette use</b> (Smoked a cigarette on at least 1 day during the 30 days before the survey.)	2.1 (1.9-2.3)
<b>Tried to quit smoking cigarettes</b> (During the past 12 months of those who smoked cigarettes.)	22.4 (21.0-23.8)
<b>Current cigar use</b> (Smoked a cigar, cigarillo, little cigar, or flavored cigar on at least one day during the 30 days before the survey.)	4.5 (4.2-4.9)
<b>Current hookah use</b> (Smoked hookah on at least 1 day during the 30 days before the survey.)	2.9 (2.7-3.1)
<b>Used first tobacco product before age 11 years</b> (Including cigarettes, cigars, little cigars, flavored cigars, hookah, and chewing tobacco.)	3.7 (3.4-4.0)
<b>Current electronic vapor product use</b> (Used at least 1 day during the 30 days before the survey.)	5.4 (5.1-5.6)

## Regional Prevalence

The table below represents tobacco use in the six regions of Cuyahoga County. This division was conducted to provide a geographic representation of the prevalence of different risk behaviors in Cuyahoga County. In 2014, we were able to weight to 5 of 6 regions in Cuyahoga County. The column under Outer Ring—West does not include data because school participation within this region was insufficient to permit weighting.

Risk Behavior	CMSD East % (95% CI)	CMSD West % (95% CI)	Inner Ring East % (95% CI)	Inner Ring West % (95% CI)	Outer Ring East % (95% CI)	Outer Ring West % (95% CI)
<b>Ever smoked cigarettes</b> (Smoked a cigarette, even one or two puffs, on at least 1 day during their life.)	12.7% (10.7-15.1)	14.3% (12.6-16.2)	8.8% (8.4-9.3)	11.3% (10.6-12.0)	3.9% (3.0-5.1)	N/A
<b>Current cigarette use</b> (Smoked a cigarette on at least 1 day during the 30 days before the survey.)	3.1% (2.2-4.2)	3.9% (3.0-5.0)	2.5% (2.3-2.7)	2.9% (2.5-3.2)	0.9% (0.7-1.3)	N/A
<b>Tried to quit smoking cigarettes</b> (During the past 12 months of those who smoked cigarettes.)	28.3% (23.7-33.4)	29.2% (23.3-35.9)	20.3% (19.0-21.5)	29.4% (28.1-30.7)	8.9% (6.9-11.5)	N/A
<b>Current cigar use</b> (Smoked a cigar, cigarillo, little cigar, or flavored cigar on at least one day during the 30 days before the survey.)	11.2% (9.1-13.8)	8.1% (6.6-9.9)	7.1% (6.7-7.5)	3.1% (2.9-3.3)	1.7% (1.3-2.3)	N/A
<b>Current hookah use</b> (Smoked hookah on at least 1 day during the 30 days before the survey.)	4.8% (3.6-6.3)	5.8% (4.5-7.5)	3.1% (2.9-3.3)	4.0% (3.7-4.2)	2.2% (1.7-2.8)	N/A
<b>Used first tobacco product before age 11 years</b> (Including cigarettes, cigars, little cigars, flavored cigars, hookah, and chewing tobacco.)	7.7% (6.0-9.9)	7.4% (6.0-9.0)	4.2% (3.9-4.5)	3.8% (3.5-4.0)	1.7% (1.3-2.1)	N/A
<b>Current electronic vapor product use</b> (Used at least 1 day during the 30 days before the survey.)	7.4% (6.1-8.8)	9.0% (7.6-10.7)	4.7% (4.5-4.9)	8.9% (8.4-9.4)	3.7% (3.2-4.3)	N/A

## Demographic Prevalence

The tables below allow for further comparisons of tobacco use between demographic groups. Data are presented by gender, race/ethnicity, grade level, and family affluence (a proxy for socioeconomic status). A statistically significant difference exists between groups if the 95% confidence intervals do not overlap.

Ever smoked cigarettes			
Category	%	CI	
<b>Gender</b>			
Female	8.8	8.2	- 9.4
Male	6.9	6.5	- 7.3
<b>Race/Ethnicity</b>			
White	5.3	5.0	- 5.5
Black	10.2	9.4	- 11.1
Hispanic	16.7	14.3	- 19.5
Other/Multiple	8.6	7.8	- 9.5
<b>Grade</b>			
7th	7.0	6.5	- 7.6
8th	8.3	7.7	- 8.9
<b>Total</b>	<b>7.8</b>	<b>7.4</b>	<b>- 8.2</b>

Current cigarette use			
Category	%	CI	
<b>Gender</b>			
Female	2.3	2.1	- 2.6
Male	1.9	1.6	- 2.1
<b>Race/Ethnicity</b>			
White	1.4	1.3	- 1.5
Black	2.4	2.0	- 2.7
Hispanic	6.8	5.4	- 8.6
Other/Multiple	2.7	2.1	- 3.5
<b>Grade</b>			
7th	1.9	1.6	- 2.1
8th	2.0	1.8	- 2.3
<b>Total</b>	<b>2.1</b>	<b>1.9</b>	<b>- 2.3</b>

Tried to quit smoking cigarettes		
Category	%	CI
<b>Gender</b>		
Female	24.9	23.2 - 26.8
Male	20.3	18.4 - 22.3
<b>Race/Ethnicity</b>		
White	21.7	20.3 - 23.3
Black	21.3	19.2 - 23.7
Hispanic	27.8	21.8 - 34.7
Other/Multiple	24.8	21.1 - 28.8
<b>Grade</b>		
7th	23.5	21.1 - 26.0
8th	21.3	19.7 - 23.0
<b>Total</b>	<b>22.4</b>	<b>21.0 - 23.8</b>

Current cigar use		
Category	%	CI
<b>Gender</b>		
Female	5.1	4.6 - 5.6
Male	4.0	3.7 - 4.4
<b>Race/Ethnicity</b>		
White	1.4	1.2 - 1.5
Black	8.5	7.7 - 9.4
Hispanic	9.9	8.3 - 11.8
Other/Multiple	5.0	3.9 - 6.3
<b>Grade</b>		
7th	3.8	3.4 - 4.4
8th	4.9	4.4 - 5.4
<b>Total</b>	<b>4.5</b>	<b>4.2 - 4.9</b>

Current hookah use		
Category	%	CI
<b>Gender</b>		
Female	3.2	2.9 - 3.5
Male	2.6	2.3 - 3.0
<b>Race/Ethnicity</b>		
White	1.5	1.4 - 1.6
Black	3.8	3.4 - 4.3
Hispanic	9.6	7.8 - 11.6
Other/Multiple	4.8	3.8 - 6.0
<b>Grade</b>		
7th	2.4	2.1 - 2.8
8th	3.0	2.7 - 3.4
<b>Total</b>	<b>2.9</b>	<b>2.7 - 3.1</b>

Used first tobacco product before age 11 years		
Category	%	CI
<b>Gender</b>		
Female	3.0	2.6 - 3.3
Male	4.4	4.0 - 4.8
<b>Race/Ethnicity</b>		
White	1.8	1.7 - 2.0
Black	5.8	5.1 - 6.5
Hispanic	7.6	6.3 - 9.3
Other/Multiple	5.1	4.8 - 5.5
<b>Grade</b>		
7th	3.5	3.2 - 4.0
8th	3.4	3.1 - 3.8
<b>Total</b>	<b>3.7</b>	<b>3.4 - 4.0</b>

Current electronic vapor product use			
Category	%	CI	
<b>Gender</b>			
Female	5.2	4.9	5.5
Male	5.5	5.2	5.9
<b>Race/Ethnicity</b>			
White	4.2	4.0	4.3
Black	5.7	5.2	6.2
Hispanic	12.9	10.8	15.4
Other/Multiple	10.0	8.9	11.2
<b>Grade</b>			
7th	5.0	4.6	5.4
8th	5.4	5.0	5.8
<b>Total</b>	<b>5.4</b>	<b>5.1</b>	<b>5.6</b>

## Cuyahoga County Trend Data

The following table presents the prevalence of tobacco use behaviors and attitudes among Cuyahoga County middle school students in 2008, 2010, 2012, 2014 and 2016. The prevalence for each year is given (when available), with 95% confidence intervals below.<sup>1</sup>

Cuyahoga County, 2008	Cuyahoga County, 2010	Cuyahoga County, 2012	Cuyahoga County, 2014	Cuyahoga County, 2016
<b>Ever smoked cigarettes</b> (Smoked a cigarette, even one or two puffs, on at least 1 day during their life.)				
21.2% (20.0-22.4)	11.9% (10.8-13.0)	8.8% (8.1-9.5)	11.9% (11.2-12.6)	7.8% (7.4-8.2)
<b>*Smoked a whole cigarette before age 11 years</b>				
4.0% (3.2-4.9)	3.4% (2.8-3.9)	2.5% (2.2-2.9)	N/A	N/A
<b>*Used first tobacco product before age 11 years</b>				
N/A	N/A	N/A	4.6% (4.2-5.0)	3.7% (3.4-4.0)
<b>Current cigarette use</b> (Smoked a cigarette on at least 1 day during the 30 days before the survey.)				
5.4% (4.8-6.0)	4.2% (3.7-4.7)	3.7% (3.3-4.0)	3.3% (3.0-3.6)	2.1% (1.9-2.3)
<b>Current cigar use</b> (Smoked a cigar, cigarillo, little cigar, or flavored cigar on at least 1 day during the 30 days before the survey.)				
8.0% (7.3-8.7)	9.0% (8.2-9.9)	6.7% (6.1-7.3)	4.8% (4.4-5.2)	4.5% (4.2-4.9)
<b>Current hookah use</b> (Smoked hookah at least 1 day during the 30 days before the survey.)				
N/A	N/A	N/A	4.3% (3.9-4.8)	2.9% (2.7-3.1)
<b>Students believe it is very wrong for someone their age to smoke cigarettes</b>				
69.0% (67.3-70.8)	68.9% (67.2-70.6)	72.5% (71.4-73.6)	N/A	N/A
<b>Students perceive that their parents believe it is very wrong for them to smoke cigarettes</b>				
88.2% (86.5-89.9)	87.9% (86.9-89.0)	89.0% (88.3-89.8)	N/A	N/A

\*Smoked a whole cigarette before age 11 years was revised in 2016 to include all tobacco products. Both items are included in this table.

<sup>1</sup> The YRBS questions with two or more data points over 2008-2016 are included. The new 2016 questions are not included for the purpose of demonstrating trends. For those questions not asked in 2016, refer to the PRCHN archives at [www.prchn.org/asp](http://www.prchn.org/asp) for more detailed information.



## Contact Information

---

**Prepared by:**

Prevention Research Center for Healthy Neighborhoods  
Case Western Reserve University  
10900 Euclid Avenue  
Cleveland, Ohio 44106-7069  
www.prchn.org

**For more information please contact:**

Erika S. Trapl, PhD  
Assistant Professor, Department of Epidemiology and Biostatistics  
Associate Director, Prevention Research Center for Healthy Neighborhoods  
Case Western Reserve University  
Erika.trapl@case.edu  
216-368-0098

Jean L. Frank, MPH  
Manager of School-Based Surveillance and Evaluation  
Prevention Research Center for Healthy Neighborhoods  
Case Western Reserve University  
Jean.frank@case.edu  
216-368-5913

Audrey E. Kinsella, MPH  
Community & Evaluation Project Coordinator  
Prevention Research Center for Healthy Neighborhoods  
Case Western Reserve University  
Audrey.kinsella@case.edu  
216-368-5746

## Partners & Funding

---

This work is funded either in whole or in part by a grant awarded by the Ohio Department of Health, Bureau of Child and Family Health Services, Child and Family Health Services Program as a sub-award of a grant issued by the Health Resources and Services Administration (HRSA) under the Maternal and Child Health Block Grant, grant award number B04MC26688, and CFDA number 93.994 and Am. Sub. H.B. 59, and by Cooperative Agreement #U87PS004165-02 from the U.S. Centers for Disease Control and Prevention, Division of Adolescent School Health (CDC-DASH).

This work was also funded by Cooperative Agreement Number 1-U48-DP-005030 from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the PRCHN and do not necessarily represent the official views of the CDC.

This work was approved by the Institutional Review Board at Case Western Reserve University, Protocol #2012-274.



CUYAHOGA COUNTY  
BOARD OF HEALTH

- 
- <sup>i</sup> U.S. Department of Health and Human Services. 2004. *The Health Consequences of Smoking: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- <sup>ii</sup> U.S. Department of Health and Human Services. 1994. *Preventing Tobacco Use among Young People: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- <sup>iii</sup> World Health Organization (Tobacco Free Initiative). *Waterpipe tobacco smoking: health effects, research needs and recommended actions by regulators* (Advisory Note). Accessed on September 17, 2013 from [http://www.who.int/tobacco/global\\_interaction/tobreg/Waterpipe%20recommendation\\_Final.pdf](http://www.who.int/tobacco/global_interaction/tobreg/Waterpipe%20recommendation_Final.pdf)
- <sup>iv</sup> U.S. Department of Health and Human Services. 1994. *Preventing Tobacco Use among Young People: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.